



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Proof by G. E. WAHLIN, Urbana, Ill.

$|1 - \sqrt{3}| < 1$. Hence (1) $0 < (1 - \sqrt{3})^{2n} < 1$.

But $(1 + \sqrt{3})^{2n} + (1 - \sqrt{3})^{2n} = E$, a rational integer, and since (1) is true, this is the integer next above $(1 + \sqrt{3})^{2n}$.

Since $\frac{(1 \pm \sqrt{3})^2}{2} = 2 \pm \sqrt{3}$, we have $\frac{(1 \pm \sqrt{3})^{2n}}{2^n} = (2 \pm \sqrt{3})^n = P \pm Q\sqrt{3}$, P and Q rational integers.

Then $\frac{(1 + \sqrt{3})^{2n}}{2^n} + \frac{(1 - \sqrt{3})^{2n}}{2^n} = \frac{E}{2^n} = 2P$. Therefore, $\frac{E}{2^{n+1}} = P$, and hence E is divisible by 2^{n+1} .

Also solved by J. Scheffer, G. B. M. Zerr, and V. M. Spunar.

PROBLEMS FOR SOLUTION.

ALGEBRA.

321. Proposed by C. C. BLAND, Attorney at Law, Rolla, Mo.

A corporation is capitalized for \$20,000. 125 shares of the par value of \$100 per share has been issued. A has 27 $\frac{19}{78}$ shares. B, C, D, E and F each have 19 $\frac{43}{78}$ shares. It is the wish of the corporation to cancel the certificates held by A, B, C, D, E, and F, and to issue new certificates to each of them in lieu of those now held by them, and to avoid the issuance of any certificate for a fraction of a share. How many shares should each receive, the whole not to exceed 200, at the same time maintaining the present interest of each in the corporation?

322. Proposed by THEODORE L. DeLAND, Treasury Department, Washington, D. C.

Take six consecutive prime numbers, as 53, 59, 61, 67, 71, and 73, and find the least whole number such that if it be divided by 59 the remainder will be 53, if it be divided by 67 the remainder will be 61, and if it be divided by 73 the remainder will be 71, and show that this least whole number and the succeeding consecutive whole numbers that will fulfill this condition as to divisions and remainders are in arithmetical progression; and also show whether or not this is a general law for n consecutive prime numbers; and if there be such a general law whether or not that general law will lead to a general law for the finding of prime numbers.

323. Proposed by E. B. ESCOTT, Ann Arbor, Mich.

Show that the relation $(a^2 + b^2 + c^2)(b^2 + c^2 + d^2) = (ab + bc + cd)^2$ can hold for real numbers only when they are in proportion.

GEOMETRY.

348. Proposed by W. J. GREENSTREET, M. A., Marling School, Stroud, England.

Two parabolas and a rectangular hyperbola circumscribe a given quadrilateral. Find a relation between the squares of the latera recta of the parabolas and the squares of the perpendiculars from the center of the hyperbola to the axes of the parabolas.